


## TREATMENT OF ANAPHYLAXIS TEST

After finishing the anaphylaxis education package please complete the following test.

1. Place each appropriate clinical feature under the adverse event to which it applies:

ANAPHYLAXIS	VASOVAGAL EPISODE
Rapid onset, usually within 5 minutes but can occur within hours	Strong central pulse
Weak/absent central pulse	Hypotension alleviated by supine or head –low position
Rash, urticaria	Normal respirations
Swelling of lips, tongue, or face	Skin pale, cool and clammy
Cough, stridor, wheeze, tachypnoea	

- a. Rapid onset, usually within 5 minutes but can occur within hours
- b. Strong central pulse
- c. Hypotension alleviated by supine or head –low position
- d. Weak/absent central pulse
- e. Rash, urticaria
- f. Swelling of lips, tongue, or face
- g. Normal respirations
- h. Skin pale, cool and clammy
- i. Cough, stridor, wheeze, tachypnoea



Score: ..... / 9

2. Treatment of anaphylaxis includes:  
(Circle correct answers)


- a. Airway maintenance
- b. O<sub>2</sub> administration by face mask at high flow rate
- c. Run for help
- d. If absent central pulse / respirations commence CPR
- e. If skin flushed or itchy commence Adrenaline injections
- f. Repeat Adrenaline injections every 2 minutes until improvement occurs
- g. The “rule of thumb” adult dose of Adrenaline 1:1000 is 5 ml
- h. The “rule of thumb” adult dose of Adrenaline 1:1000 is 0.5 ml
- i. Repeat Adrenaline injections every 5 minutes until improvement occurs
- j. Monitor for at least 4 hours following recovery

Score: ..... / 6

**3. Most life-threatening adverse events begin:**

(Circle correct answer)

- a. Within 10 minutes of vaccination
- b. Within 4 hours of vaccination
- c. Within 7 days of vaccination



**4. Recipients of vaccines should:**


(Circle correct answer)

- a. Leave as soon as the vaccine has been given
- b. Remain in the vicinity of the place of vaccination for at least 15 minutes
- c. Stay for up to two hours after administration
- d. Stay for 30 minutes after vaccination

**5. Adrenaline should be administered via:**

(Circle correct answer)

- a. Intravenous injection
- b. Sub cutaneous injection
- c. Orally
- d. Deep intra muscular injection



**6. Adult adrenaline is given as:**

(Circle correct answer)

- a. Adrenaline 1:5000
- b. Adrenaline 1:10 000
- c. Adrenaline 1:1000
- d. Adrenaline 2:1000

**7. Adult dose of Adrenaline 1:1000 is:**

(Circle correct answer)

- a. 5.0mL
- b. 0.05mL
- c. 0.5mL

Score: ..... / 5

Date: .....

Total score - ..... / 20

Participant: .....

Supervisor: .....

Signed: .....

Signed: .....